

Certificate of Analysis

Catalog Number	BP63038
Product Name	Anti-NANOG antibody

Physical and Chemical Properties

Molecular Weight	35-40 kDa
GenBank	BC160187
Uniprot	Q9H9S0
Concentration	660 μg/ml
Form	Liquid
Storage Instruction	10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at -20°C. Do Not Aliquot.
Chemical Structure OR Tested Image	250 - 150 - 100 - 75 - 50 - 37 - NANOG 25 - 15 - 10 -

Product Information

Description	Nanog is a member of the homeobox family of DNA binding transcription factors and has been shown to maintain embryonic stem (ES) cell self-renewal independently of leukemia inhibitory factor (LIF)/Stat3. Nanog mRNA is present in pluripotent mouse and human cell lines, and absent from differentiated cells. Functionally, Nanog works together with other key pluripotent factors (Oct4, Sox2, and Lin28) to reprogram human fibroblasts and generate induced pluripotent stem (iPS) cells. These key factors form a regulatory network to support or limit each other's expression level, which maintains the properties of ES cells. Affinity purified rabbit anti-Nanog can be used to demonstrate pluripotency of ES and IPS cells. There are two kinds of variants could recognized by NANOG, one is normal form (~39kd), the other is post-translation modified form (~48kd) (21136380). Nanog exists two isoforms with molecular weight 34.4 kDa and 31.9 kDa.
Tested Applications	WB: 1:1000; IF: 1:100-1:300; IHC:1:50-1:200
Species Reactivity	Human, Mouse, Rat
Host Species/Isotype	Rabbit/IgG

Analytical Data

Quality Assurance	The biological and chemical parameters such as concentration, purity, application and specificity of the tested antibody comply with the above-mentioned criteria of the product.
-------------------	---

Purdue Bioscience Inc.

750 50th St, Brooklyn, NY 11220, USA

http://www.purduebio.com

1-877.618.7311

info@purduebio.com

v2 Revision on 12/28/2022